

Lawrence Dental Group

500 3rd Ave, Kingston, PA 18704

570-288-5588

OPALESCENCE INSTRUCTIONS FOR AT HOME WHITENING

Although take-home whitening is a relatively new procedure, it has been effective and safe for most patients. There may be some patients for whom whitening is either not effective or possibly not the most preferable treatment. The amount of whitening that you obtain during the procedure is dependent on four factors:

1. The length of time in each 24-hour period the tray is worn.
2. The number of sessions the tray is worn.
3. The susceptibility of your teeth and/or internal colors to the whitening agent.
4. Habits you have that discolor teeth: smoking, coffee and/or tea drinking, red or dark wines, etc.

PROCEDURE

Depending upon the strength of solution you are given, the amount of time for wear is as follows:

OPALESCENCE 20% 2-4 HOURS PER DAY

OPALESCENCE 35% 1/2 HOUR PER DAY

DO NOT SLEEP WITH BLEACH/TRAYS IN YOUR MOUTH.

PROCEDURE

1. Before inserting trays, brush and floss teeth thoroughly. Also, brush the custom trays so they are clean. Dry the trays. Be sure not to flex the tray too much while cleaning and do not use hot water.
2. Load the tray with the whitening gel. Express the gel evenly into the deepest, outer most portion of the trays. Use a very small amount. **A large amount is not needed.**
3. Seat the trays completely onto the teeth.
4. Gently press the tray with a clean finger to adapt the soft tray material against the teeth on the inside (tongue side) and the outside (lip side) edges of the tooth/gum area. Pressing too firmly will express too much gel out of the tray.
5. Wipe off any excess gel with a clean finger or a dry toothbrush.
6. Do not disturb the trays when wearing by lifting with tongue, fingers, etc. Take care not to bite with pressure on the tray. This may cause excess solution to seep out and set on gum tissue resulting in a tissue burn.

7. Remove trays after wearing the appropriate amount of time. Brush teeth thoroughly with toothpaste. Rinse twice; do not swallow rinsed gel. Brush tray gently with soft brush and rinse with cool water.

8. On the average treatment should take 5-7 days depending on patient's goals, stains, and comfort.

9. The 35% is a very strong strength of whitening. Both whitening types have carbamide peroxide along with potassium nitrate and fluoride to strengthen enamel and decrease sensitivity.

POSSIBLE SIDE EFFECTS

Many times, patients will experience increased sensitivity to cold during treatment. Before, during and after whitening treatment, you can use a sensitivity toothpaste. When using this type of toothpaste, floss, brush w/the toothpaste, spit, but do not rinse with anything (water/mouthwash) after brushing.

Some patients have reported temporary discomfort during whitening, such as gum and/or tooth sensitivity, tongue and lip soreness, throat irritation, or moderate, continuous teeth pain. Acidic citric foods may increase sensitivity temporarily.

If any of these symptoms occur and are more than mild, are persistent, or if you have any questions or concerns, call our office. These side effects most always resolve in 1-3 days after interruption or completion of treatment. **If gum tissue becomes whitish, please discontinue whitening treatment until gum tissue resumes normal color.**

PRECAUTIONS

Avoid dark foods or drinks that may stain for 24 hours after whitening.
For example: Coffee, tea, red wine, tobacco, dark berries

Keep and store the whitening agent out of heat or direct sunlight at all times as the bleaching agent will chemically break down. You may store extra material in the refrigerator. **NEVER** use any household or commercial bleaching agents in your mouth!

Regular dental checkups and cleanings are important before and after bleaching to maintain a healthy smile.